




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : B01D 15/00	A1	(11) International Publication Number: WO 89/01813 (43) International Publication Date: 9 March 1989 (09.03.89)
(21) International Application Number: PCT/US88/02989 (22) International Filing Date: 29 August 1988 (29.08.88) (31) Priority Application Number: 091,728 (32) Priority Date: 1 September 1987 (01.09.87) (33) Priority Country: US (71) Applicant: PROCESS BIOTECHNOLOGY, INCORPORATED [US/US]; 1215 Bardstown Trail, Ann Arbor, MI 48105 (US). (72) Inventors: WANG, Henry, Y. ; 2388 Bunker Hill Road, Ann Arbor, MI 48105 (US). TSAO, I-Fu ; 24 Grandin Drive, Flemington, NJ 08822 (US). (74) Agents: HOFER, Ronald, L. et al.; Harness, Dickey & Pierce, 1500 North Woodward Avenue, Birmingham, MI 48011 (US).		(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published With international search report.

(54) Title: CELLULAR RECEPTOR ADSORBENTS FOR REMOVAL OF VIRAL CONTAMINANTS FROM BLOOD AND COMPLEX PROTEIN SOLUTIONS

(57) Abstract

Cellular receptor adsorbents (10) for removing viral contaminants from whole blood and protein solutions are prepared by immobilizing and stabilizing cells or portions thereof (12) bearing cellular receptors (14) for the target contaminants (18). Removal of the viral contaminants from solution is effected by affinity binding (24) of viral proteins of the contaminant and their corresponding immobilized cellular receptors.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CF Central African Republic		SE Sweden
CN China		SN Senegal
CO Colombia		
CR Costa Rica		
CZ Czechoslovakia		
DE Germany	MC Monaco	DJ Djibouti
DK Denmark	MG Madagascar	
FI Finland		

AT Austria
AU Australia
BB Barbados
BE Belgium
BG Bulgaria
BJ Benin
BR Brazil
CF Central African Republic

FR	France
GA	Gabon
GB	United Kingdom
HU	Hungary
IT	Italy
JP	Japan
KP	Democratic People's Republic of Korea

ML	Mali
MR	Mauritania
MW	Malawi
NL	Netherlands
NO	Norway
RO	Romania
SD	Sudan
SE	Sweden
SN	Senegal

DK Denmark
FI Finland

MC Monaco
MG Madagascar

- 1 -

CELLULAR RECEPTOR ADSORBENTS FOR REMOVAL OF VIRAL
CONTAMINANTS FROM BLOOD AND COMPLEX PROTEIN SOLUTIONS

BACKGROUND OF THE INVENTION

The present invention relates generally to the removal of
5 viruses from solution and, more particularly, to the preparation of
immobilized cellular receptor adsorbents used for removal of viral
contaminants from blood, plasma, serum, and blood-derived products.

Viruses in blood products used for intravenous
administration present well-recognized risks of infection despite
10 health history evaluations and serological screening. Infections
which have been transmitted in this manner include hepatitis and AIDS.
Similarly, the presence of viruses in animal sera supplied for cell
culture often results in contamination of the culture.

Numerous attempts have been made to prevent or eliminate
15 viral contamination in solution by inactivating the viruses. However,
the problems encountered in viral inactivation in blood and protein
solutions are distinct from the problems of viral inactivation in
other aqueous solutions. Viruses in many aqueous solutions can easily
be inactivated by pasteurization, radiation and/or chemical
20 sterilization. In contrast, whole blood, some blood protein fractions
and most protein solutions such as protein-based pharmaceuticals are
sensitive to such physical or chemical treatments.

Pasteurization or thermal inactivation, a widely accepted
method of sterilizing certain proteins such as albumin, commonly
25 results in a marked reduction or elimination of functional activity of
labile proteins such as clotting factors. Use of high concentrations
of polysaccharides to stabilize proteins during pasteurization
unfortunately also increases the thermoresistance of the viruses.
Viral inactivation by treatment of serum or plasma with
30 beta-propiolactone and UV irradiation and by lipid solvent treatments
results in a significant loss of functional activity of proteins.

Loss of protein functional activity is similarly encountered
when blood-based solutions are treated by hydrophobic interaction
chromatography to remove the viruses from solution. Antibody-based
35 chromatography using polyclonal antibodies and monoclonal IgM as
affinity ligands preserves protein activity, but has the disadvantage

- 2 -

of being too specific. The presence of mutants and different types of antigenic determinants thus often leads to failure of binding the target viruses. Removal of viruses from solution by filtration is also not practical because of the trade-off between required pore size restriction and the low filtrate flux.

Thus, there is a need for an improved method for removal of viral contaminants especially from blood and complex protein solutions. The present invention provides such an improved method and an adsorbent for carrying out the method. Further understanding of the invention will be had from the following disclosure taken in conjunction with the drawing and claims.

SUMMARY OF THE INVENTION

The present invention provides a novel approach for removal of viral contaminants such as viruses and virus-infected cells in which immobilized cellular receptor adsorbents prepared in accordance with the method of the present invention are used as affinity ligands to bind and remove viral contaminants from solution. In accordance with the present invention, immobilized cellular receptor adsorbents for a target viral contaminant are prepared by a method comprising the steps of:

(A) selecting a cell bearing a cellular receptor for said target viral contaminant;

(B) immobilizing at least a portion of said cell, said portion bearing said cellular receptor thereon, onto a biologically-compatible support surface; and

(C) modifying said cell portion to stabilize said portion.

The present invention thus provides a means for sterilization of blood or other protein solutions by removal of viral contaminants by cellular receptor adsorbents prepared by the method of the invention. Sterilization by cellular receptor adsorbents of the present invention offers several advantages over conventional methods of sterilization. Since viral protein-cellular receptor interactions are specific bindings with high affinity, product loss due to non-specific adsorption is reduced to a minimum. Also, cellular receptors possess a much broader adsorption spectrum than

- 3 -

immunoabsorbents, thus widening the range of viruses removed to include variants of the targeted contaminants.

A further advantage of the present invention is that use of cellular receptor adsorbents of the present invention does not result in a significant decrease of protein function or therapeutic effectiveness of the solution being processed. Also, lysis of whole cells such as blood cells can be kept to a minimum. Thus, whole blood can be easily and effectively sterilized by the adsorbents of this invention. Additionally, cellular receptor adsorbents of the present invention can be used not only to remove whole viruses, but other viral contaminants such as infected cells carrying viral-encoded proteins on their surface. Further, cellular receptor adsorbents of the present invention are also relatively inexpensive and easy to prepare.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic diagram illustrating a preferred embodiment of the method of virus adsorption using an immobilized cellular receptor adsorbent of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, cellular receptor adsorbents for removing viral contaminants from solution are prepared by immobilizing and stabilizing cells or cell parts bearing cellular receptors on a biologically-compatible support surface. The term viral contaminants, as used herein, includes whole viruses, viral fragments, infected cells expressing viral proteins or any other contaminants which contain or mimic viral proteins to which cellular receptors exhibit binding affinity.

Referring now to Figure 1, a cellular receptor adsorbent of the present invention is illustrated and indicated generally by the numeral 10. Cellular receptor adsorbent 10 generally comprises a cell or cell part 12 bearing at least one cellular receptor 14 thereon which has been immobilized on a support surface 16 and thereafter stabilized to prevent disruption of that adsorbent. Cellular receptor 14 or cell part 12 to be immobilized is selected for specificity for a target virus 18.

- 4 -

In use, cellular receptor adsorbent 10 is brought into contact with a solution contaminated with target virus 18. As depicted in Figure 1A, the solution may be a complex protein solution containing a plurality of proteins 20, 22 in addition to target virus 18. As shown in Figure 1B, after contact with cellular receptor adsorbent 10, target virus 18 in solution will bind to a cellular receptor 14 of the adsorbent, as shown at 24, while other proteins 20, 22 stay in solution. The protein solution, now substantially free of target virus 18 which is bound to cellular receptor adsorbent 10, can then be separated from adsorbent, as illustrated in Figure 1C, and collected for further use.

Cells to be immobilized in accordance with the present invention are selected for their membrane receptor specificity for the viral contaminants to be removed from solution. Proteins of cells, such as cell ghosts or purified plasma membrane, may also be used in conjunction with or in lieu of whole cells. Additionally, more than one type of cell or portion thereof with overlapping or different viral specificities may be selected to further extend the adsorption spectrum. Table 1 sets forth examples of "receptor maps" of certain cells.

TABLE 1.: EXAMPLES OF RECEPTOR MAPS OF VIRUSES

<u>Cell</u>	<u>Virus</u>
T4-lymphocyte	HTLV-I, HTLV-II, HTLV-III (HIV)
Hepatocyte	Hepatitis B v. Non-A, non-B hepatitis v.
Human foreskin fibroblasts	Human cytomegalovirus
Rat T-lymphocyte	Influenza B(H2N2), A(H2N2), A(H3N2) Newcastle disease v. Sendai v.

- 5 -

	HeLa	Human Rhinovirus types 2,1A,1B
		Human Rhinovirus types 14,3,5,15,39,41,51
5		Polioviruses Coxsackieviruses Adenoviruses
	Erythrocyte	New Castle Dis. v. Influenza A Sendai v. Rubella v. Reovirus
10		
	Mouse A-9 FLC	MVM
15	GMK AH-1 Vero SIRC HeLa S3 HEP-2	HSV-1 HSV-2
20	HEL	
	Murine teratoma F9, F12, PCC4 Mouse embryo fibroblast Daudi cells	Coxsackie B3 Mouse hepatitis Sindbis Semliki Forest Lymphocytic choriomeningitis Pichinde Vesicular stomatitis HSV-1 Cytomegalovirus
25		
30	BHK-21	FMDV types A ₁₂ , O _{1B} , C _{3res}

- 6 -

	Murine & Human Ependymal cell	Reovirus type 1 Mumps v. Parainfluenza type 3 HSV-1
5	SEM FLC L-929 HeLa	Mengovirus EMCV
10	L ₈ CL3-U	Coxsackie v. types A2,A5,B3 HSV-1
	HeLa L cell	Phinovirus 24 serotypes
15	BHK-21 Neuroblastoma C1300	ERA Rabi v. RV 194-2 VSV Reovirus type 3 West Nile v.

20 With the advent of genetic engineering, selected cells can also be quantitatively programmed to over-express certain receptor proteins on their surface to maximize their adsorption capacity for a specific virus. Alternatively, a cell line with an expanded adsorption spectrum or multispecificity for several viruses may be constructed by incorporating genes encoding the corresponding cellular

25 receptors.

The cells or cell parts to be immobilized in accordance with the method of the invention can be either anchorage-dependent or anchorage-independent. It should be appreciated that the nature and adhesive properties of the cell surface can be affected by its

30 composition, charge, and age. The composition, charge, and shape of the support material also influences the attachment of cells to the

- 7 -

Materials used for the cell support surface can be beads, membranes or other conventional supports which are biologically-compatible with the biological systems involved. By biologically-compatible support surface is meant a support surface which does not substantially adversely affect, or release any undesired components or products which substantially adversely affect, the cellular receptor adsorbents, the immobilization or stabilization processes, or the solution being processed. Suitable materials for support surfaces include polymeric materials, for example polysaccharides such as alginate, agarose, carrageenan, dextran, and cellulose, as well as gelatin, glass, acrylic copolymers, titanium oxide, ceramic and other compounds or materials on which cell immobilization can be achieved. Materials of a non-porous nature are preferred in order to eliminate non-specific adsorptions of desired proteins onto an intraporous surface.

A variety of surface immobilization methods can be employed to attach the selected cells or cell parts to the support surface. For example, cells can be covalently attached by coupling them to a support surface previously activated by a cross-linking agent. In ionic attachment, the support surface is treated to provide a polyanionic or polycationic surface to which cells can attach through charge interactions.

Alternatively, cells or cell parts can be immobilized by affinity binding which involves specific interactions between molecules on the cell surface and corresponding proteins immobilized on the support surface. Typical affinity interactions include antigen-antibody, hormone-receptor, and lectin-glycoprotein binding. Cell attachment can also be achieved by cultivating anchorage-dependent cells on a suitable support surface, such as microcarrier beads.

In addition to immobilizing the cells or cell parts on the support surface, the plasma membranes of the immobilized cells or parts thereof must be stabilized. Animal cells, in contrast to bacterial cells, lack cell walls which provide a rigid mechanical support against shear stress and osmolarity changes in the environment. As a result, their plasma membranes are particularly sensitive to disruption and disintegration.

- 8 -

Disruption of the immobilized plasma membrane of the adsorbent can destabilize the surface for viral protein binding and also contaminate the solution being processed by the release of cell membrane components into the solution. If whole cells are being used, membrane disruption can also lead to cell leakage of intracellular components into and further contamination of the solution. Stabilization of the plasma membrane is thus necessary to prevent such contamination. Additionally, intercellular, as well as intracellular, stabilization of membrane proteins will further enhance stability of the membrane surface and will provide a more contiguous binding surface for viral contaminants.

Plasma membranes can be stabilized by treatment with a cross-linking agent to cross-link the membrane proteins. Suitable cross-linking agents include glutaraldehyde and other aldehydes, benzoquinone, carbonyldiimidazole, periodate and the like. The toxicity of the cross-linking agent employed should be as low as possible, and should preferably have a minimal effect on glycoproteins since most known cellular receptors for viruses are glycoproteins. The stabilization of the plasma membranes by the cross-linking agent should also not significantly hamper the functional activities of the cellular receptors. This can be achieved by selecting a suitable cross-linking agent at an appropriate concentration which is generally very low, for example, 0.1% by volume for glutaraldehyde.

Alternatively, the risk of contamination may be minimized by preparing adsorbents from receptor-bearing cell parts or cells whose intracellular components have been released beforehand in a controlled manner. For example, immobilized eukaryotic cells can be hypotonically lysed to release their cell contents without unduly affecting their membrane composition. Subsequent stabilization of the immobilized membranes will then provide a more stable and contiguous surface for viral binding.

In use, the immobilized and stabilized adsorbent is contacted with whole blood or any other solution to be processed for a sufficient time to permit the adsorbent to bind the target viral contaminant and remove substantially all contaminants from solution. By substantially all is meant the reduction of virus or viral contaminant to a level below detection by conventional methods such as

- 9 -

hemagglutination and plaque assays. Optionally, after processing of the solution has been completed and the solution collected, the cellular receptor adsorbent is disposed of to eliminate any risk of release of undesired materials or contaminants into a second solution during re-use of the adsorbent.

It will be appreciated that, although cellular receptor adsorbents of the present invention will remove target viral contaminants from a contaminated solution, the adsorbents of the invention can also be used to routinely treat blood or other solutions not specifically known to be contaminated. Such routine processing would be highly desirable for whole blood and blood products prior to intravenous administration to reduce the risk of infection of the recipient.

In Example 1, discussed in detail below, virus removal by a cellular receptor adsorbent prepared in accordance with the method of the present invention is exemplified by the adsorption of influenza A virus by immobilized chicken erythrocyte membranes. Chicken erythrocytes with cellular receptors for influenza A virus were first immobilized on polycationic beads in a column. Ghosts of the immobilized erythrocytes were then prepared by subjecting them to hypotonic conditions and the membranes were then resealed. The immobilized erythrocyte membranes were stabilized by treatment with low concentrations of glutaraldehyde. Influenza A virus in buffered solution was twice passed through the column of immobilized erythrocyte membranes to effect removal of substantially all of the influenza A viral population from solution.

In Example 2, a cellular receptor adsorbent for cytomegalovirus removal was prepared by cultivating anchorage-dependent human foreskin fibroblasts on microcarriers and then stabilizing the cell membranes with low concentrations of glutaraldehyde. In Example 3, a cellular receptor adsorbent for removal of herpes simplex virus was identically prepared by cultivating anchorage-dependent BSC-1 cells on microcarriers and stabilizing them with low concentrations of glutaraldehyde. Similar cellular receptor adsorbents can be prepared for removal of hepatitis viruses using anchorage-dependent hepatocytes.

- 10 -

In Example 4, cellular receptor adsorbent for removal of human T-cell lymphotropic viruses (HTLV) was prepared by affinity binding T-cells to carboxylate modified latex microspheres bearing a lectin specific for human T cells. The T-cell membranes were then stabilized by low concentrations of glutaraldehyde.

It will be appreciated that the method of the present invention is versatile and subject to variation and that the specific cellular receptor adsorbent and its preparation will be dependent upon the type of cells, immobilization supports and target contaminants involved. All percentages given herein are by volume unless otherwise indicated.

EXAMPLE 1

Chicken erythrocytes were washed twice with phosphate buffer saline (PBS) to remove extracellular proteins and resuspended in sucrose-acetate buffer (SAB). The PBS contained 150 mM NaCl and 5mM sucrose, pH 5.2. Polyacrylamide beads coated with polyethyleneimine were used as a polycationic solid support (Affi-Gel 731). The beads were hydrated and then washed three times with 0.2 M NaCl solution, twice with distilled water and twice with SAB. 0.5 ml of 50% bead suspension was added dropwise to a same volume of 50% erythrocyte suspension with occasional gentle shaking. All percentages given herein and in Examples 2-4 are by volume unless otherwise indicated. After the erythrocytes were bound to the surface, dextran sulfate (1 mg/ml, M.W. 5,000) in SAB was added to neutralize the surface sites not covered by the cells. The immobilized erythrocyte beads were then washed three times with SAB, gently enough to avoid cell disruption and detachment.

Hypotonic treatment to obtain ghost cells was performed in hemolysis medium (4 mM $MgSO_4$, 3.8 mM CH_3COOH , pH 3.8) for 5 minutes. 10 ml of PBS was added to resume the tonicity. The supernatant containing intracellular debris and vesicles was aspirated after the beads settled down. To reseal the erythrocyte ghosts, 10 ml of PBS was added and followed by incubation of the suspension at 37°C for 15 minutes. The beads were then washed three times with PBS. The immobilized membranes were then stabilized by incubating the beads with 0.1% glutaraldehyde at room temperature for one hour with gentle

- 11 -

shaking. Finally, the immobilized and stabilized membranes were washed three times with PBS to remove excess cross-linking agent and other materials and stored at 4°C.

Table 2 demonstrates the titer reduction of influenza A virus in minimal essential medium (MEM) with Earle's salts supplemented with 10% fetal bovine serum (FBS) after being passed through the column packed with immobilized erythrocyte beads. The passes through the column were performed at 4°C because viral neuraminidase, which could destroy the receptors for hemagglutinin and facilitate the elution of virions from cell surface, is not active in the low temperature range. As shown in Table 2, the hemagglutination (HA) tests showed that viruses were removed below detectable levels after the second pass. The HA tests of the subsequent passes were negative and indicated that the viruses were not being released from the beads.

TABLE 2. THE AFFINITY OF INFLUENZA A VIRUSES BY
SURFACE-IMMOBILIZED ERYTHROCYTE GHOSTS

HA unit	1	2	3	4	8	16	32	64	128
control	+	+	+	+	+	+	+	+	-
1st pass	+	-	-	-	-	-	-	-	-
2nd pass	-	-	-	-	-	-	-	-	-
3rd pass	-	-	-	-	-	-	-	-	-
4th pass	-	-	-	-	-	-	-	-	-
5th pass	-	-	-	-	-	-	-	-	-

* Hemagglutination tests were performed with two-fold serial dilution. The positive sign means the viruses are present and

- 12 -

preventing the settlement of erythrocytes, while the negative sign indicates the virus titer is below the detectable limit of the HA test.

EXAMPLE 2

5 Anchorage-dependent human foreskin fibroblasts were grown on microcarriers (Superbead, Flow laboratories) in minimal essential medium (MEM) with Earle's salts supplemented with 10% fetal bovine serum (FBS) and 1.1 g/l sodium bicarbonate and incubated at 37°C in a humidified 4% CO₂ - 96% air atmosphere. After cell growth reached
10 confluency, the microcarriers were washed twice using HEPES-buffered saline (HBS), then incubated with 0.05% glutaraldehyde at room temperature for one hour with occasional shaking. The residual glutaraldehyde was removed by washing the microcarrier four times using HBS. The final preparation which can be used to adsorb
15 cytomegalovirus was resuspended in HBS and stored at 4°C.

EXAMPLE 3

Anchorage-dependent BSC-1 cells were grown on microcarriers (Superbead, Flow laboratories) in minimal essential medium (MEM) with Earle's salts supplemented with 10% fetal bovine serum (FBS) and 1.1
20 g/l sodium bicarbonate and incubated at 37°C in a humidified 4% CO₂ - 96% air atmosphere. After cell growth reached confluency, the microcarriers were washed twice using HEPES-buffered saline (HBS), then incubated with 0.05% glutaraldehyde at room temperature for one hour with occasional shaking. The residual glutaraldehyde was removed
25 by washing the microcarrier four times using HBS. The final preparation which can be used to adsorb herpes simplex viruses was resuspended in HBS and stored at 4°C.

EXAMPLE 4

30 A layer of buffy coat was separated from human venous blood by equilibrium centrifugation using a Ficoll-Paque medium. Buffy coat cells, namely B cells, T cells and monocytes, were incubated with carboxylate modified latex microspheres with average diameter of 2μ at room temperature for one hour. The microspheres bear the lectin helix pomatia which specifically binds human T cells. A final

- 13 -

centrifugation concentrated and separated out a fairly pure fraction of immobilized T cells. The clusters of T cells and microspheres were incubated with 0.05% glutaraldehyde solution at room temperature for one hour with gentle shaking. The final preparation which can be used

5 to adsorb HTLV viruses was resuspended in HBS and stored at 4°C.

What is Claimed is:

1. A method of preparing a cellular receptor adsorbent for removing a target viral contaminant from a solution comprising the steps of:
 - 5 (A) selecting a cell bearing a cellular receptor for said target viral contaminant;
 - (B) immobilizing at least a portion of said cell, said portion bearing said cellular receptor thereon, onto a biologically-compatible support surface; and
 - 10 (C) modifying said cell portion to stabilize said portion.
2. The method of Claim 1, wherein step (C) is carried out by treating said cell portion with a cross-linking agent.
3. The method of Claim 1, wherein step (B) is carried out
15 by covalently binding said portion of said cell onto said biologically-compatible support surface.
4. The method of Claim 1, wherein step (B) is carried out by ionic binding of said portion of said cell onto said biologically-compatible support surface.
- 20 5. The method of Claim 1, wherein step (B) is carried out by affinity binding of said portion of said cell onto said biologically-compatible support surface.
6. The method of Claim 1, wherein said cells are anchorage-dependent cells.
- 25 7. The method of Claim 2, wherein said cross-linking agent comprises glutaraldehyde.
8. The method of Claim 2, wherein said cell portion is a portion of a membrane of said cell and said membrane is prepared by

- 15 -

9. The method of Claim 1, wherein said cellular receptor is overexpressed.

10. The method of Claim 2, wherein said target viral contaminant is selected from a group of viruses consisting of hepatitis viruses, herpes viruses, retroviruses and T-cell lymphotropic viruses.

11. The adsorbent prepared by the method of Claim 2.

12. A method of removing substantially all of a target virus population from a solution comprising the steps of:

10 (A) selecting a cell bearing a cellular receptor for said target virus population;

(B) immobilizing at least a portion of said cell, said portion bearing said cellular receptor thereon, onto a biologically-compatible support surface;

15 (C) modifying said cell portion to stabilize said portion; and

(D) contacting said solution with said immobilized stabilized cell portion for a time sufficient to permit binding of said viruses of said population to said cellular receptors.

20 13. The method of Claim 12, wherein step (C) is carried out by treating said cell portion with a cross-linking agent.

14. The method of Claim 12, wherein said solution comprises a protein solution.

25 15. The method of Claim 12, wherein said solution comprises whole blood.

16. The method of Claim 12, wherein said time sufficient to permit binding is further sufficient to reduce the titer of said target virus to a level below detection.

- 16 -

17. The method of Claim 13, wherein said cells are erythrocytes and said target virus population comprises influenza viruses.

18. The method of Claim 13, wherein said cells are BSC-1
5 and said target virus population comprises herpes simplex viruses.

19. The method of Claim 13, wherein said cells are human fibroblasts and said target virus population comprises cytomegalovirus.

20. The method of Claim 13, wherein said cells are
10 T-lymphocytes and said target virus population comprises T-cell lymphotropic viruses.

21. The method of Claim 13, wherein said cells are hepatocytes and said target virus population comprises hepatitis viruses.

22. A method for sterilizing a blood-based protein solution
15 comprising the steps of:

(A) selecting a cell having a membrane bearing cellular receptors for a target viral contaminant;

(B) immobilizing at least a portion of said membrane on
20 a biologically-compatible support surface;

(C) stabilizing said portion of said membrane by treating said portion with a cross-linking agent;

(D) contacting said stabilized immobilized portion of
said membrane with said blood-based protein solution for a time
25 sufficient to permit binding of said viral contaminant to said cellular receptors; and


(E) collecting said blood-based solution after said binding of said viral contaminant.

- 17 -

23. A method for treating a protein solution comprising the steps of:

- (A) selecting a cell having a membrane bearing a cellular receptor for a target viral contaminant;
- 5 (B) immobilizing said portion of said membrane on a biologically-compatible support surface;
- (C) modifying said portion of said membrane to stabilize said portion;
- (D) contacting said immobilized stabilized portion of
- 10 said membrane with said solution; and
- (E) collecting said solution after said contact.

24. The method of Claim 23, wherein said protein solution comprises blood.



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US88/02989

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(4): BOLD 15/00 U.S. CL.: 210/679		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	210/198.2, 500.21-500.43, 501, 502.1, 601, 635, 656, 679, 905, 908; 435/5, 176, 177, 180, 181, 239; 530/350, 380, 415, 826; 424/89, 101; 422/1, 28; 604/4, 5; 502/400, 401, 403	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	US, A, 4,361,484 (LARSSON) 30 NOVEMBER 1982 SEE THE ENTIRE DOCUMENT.	1-24
Y	US, A, 4,282,315 (LUDERER) 04 AUGUST 1981 SEE THE ENTIRE DOCUMENT.	1-24
A	US, A, 3,957,580 (NELSON) 18 MAY 1976	1-24
A	DALEN, A., "THE EFFECT OF GLUTARALDEHYDE ON THE STABILITY OF ERYTHROCYTES AND ON VIRUS RECEPTOR SUBSTANCES", ACTA. PATH. MICROBIOL. SCAND., SECT. B, VOL. 84, PAGES 196-200, (1976).	1-24
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>[*] Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
12 October 1988		01 DEC 1988
International Searching Authority		Signature of Authorized Officer